

AD-A211 545



# LIFETIME OF SINGLET OXYGEN IN AQUEOUS HYDROGEN PEROXIDE (BHP)

M.A.J. Rodgers  
M.A. Shand

Center for Photochemical Sciences  
Bowling Green State University  
Bowling Green, Ohio 43403

May 1989

Final Report

Approved for public release; distribution unlimited.

WEAPONS LABORATORY  
Air Force Systems Command  
Kirtland Air Force Base, NM 87117-6008

DTIC  
ELECTE  
AUG 21 1989  
S B D

89 8 21 199

This final report was prepared by the Center for Photochemical Sciences, Bowling Green State University, Bowling Green, Ohio under Contract F29601-87-K-0076, Job Order 33260345 with the Weapons Laboratory, Kirtland Air Force Base, New Mexico. Mr Leonard Hanko (ARDI) was the Laboratory Project Officer-in-Charge.

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely Government-related procurement, the United States Government incurs no responsibility or any obligation whatsoever. The fact that the Government may have formulated or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication, or otherwise in any manner construed as licensing the holder or any other person or corporation; or as conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.


This report has been authored by a contractor of the United States Government. Accordingly, the United States Government retains a nonexclusive, royalty-free license to publish or reproduce the material contained herein, or allow others to do so, for the United States Government purposes.

This report has been reviewed by the Public Affairs Office and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

If your address has changed, if you wish to be removed from our mailing list, or if your organization no longer employs the addressee, please notify WL/ARDI, Kirtland AFB, NM 87117-6008 to help us maintain a current mailing list.

This technical report has been reviewed and is approved for publication.

  
LEONARD HANKO  
Project Officer

  
MICHAEL T. WHITE  
Major, USAF  
Chief, Advanced Chemical Laser Br

FOR THE COMMANDER

  
ERIC J. JUMPER  
Lt Col, USAF  
Chief, Devices Division

DO NOT RETURN COPIES OF THIS REPORT UNLESS CONTRACTUAL OBLIGATIONS OR NOTICE ON A SPECIFIC DOCUMENT REQUIRES THAT IT BE RETURNED.


SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFWL-TR-89-07	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			7a. NAME OF MONITORING ORGANIZATION Weapons Laboratory	
6a. NAME OF PERFORMING ORGANIZATION Center for Photochemical Sciences		6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State, and ZIP Code) Kirtland Air Force Base, NM 87117-6008	
6c. ADDRESS (City, State, and ZIP Code) Bowling Green State University Bowling Green, Ohio 43403		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F29601-87-K-0076		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	10. SOURCE OF FUNDING NUMBERS	
8c. ADDRESS (City, State, and ZIP Code)		PROGRAM ELEMENT NO. 62601F	PROJECT NO. 3326	TASK NO. 03
		WORK UNIT ACCESSION NO. 45		
11. TITLE (Include Security Classification) LIFETIME OF SINGLET OXYGEN IN AQUEOUS HYDROGEN PEROXIDE (BHP)				
12. PERSONAL AUTHOR(S) Rodgers, M.A.J.; and Shand, M.A.				
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 18Sep87 to 22Feb89		14. DATE OF REPORT (Year, Month, Day) 1989 May
15. PAGE COUNT 26				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	Singlet delta oxygen; Deuterated basic hydrogen peroxide	
11	06	02	Lifetime	
20	13		Quenching rates; Bimolecular rate constants	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  Singlet delta oxygen lifetime in deuterated basic hydrogen peroxide was measured. The contributing bimolecular rate constants $k_{00}$ and $k_{0,2}$ were measured as a function of temperature. The measured lifetime of 20 $\mu$ s compared favorably to the lifetime of 24 $\mu$ s arrived at by using contributing bimolecular rates.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL LEONARD HANKO			22b. TELEPHONE (Include Area Code) (505) 844-0661	
			22c. OFFICE SYMBOL ARDI	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## SUMMARY

The lifetime ( $\tau_{\Delta}$ ) of singlet molecular oxygen  $O_2(^1\Delta_g)$  in aqueous solution is little affected by the presence of the species  $OH^-$ ,  $H_2O_2$  or  $OD^-$  and  $D_2O_2$ , compared with its lifetime in either  $H_2O$  or  $D_2O$ , respectively.

The bimolecular quenching rate constants for the quenching of  $O_2(^1\Delta_g)$  by  $OD^-$  and  $D_2O_2$  have been determined as  $5.5 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$  and  $8.8 \times 10^2 \text{ M}^{-1}\text{s}^{-1}$ , respectively. A direct measurement of the bimolecular quenching rate constants for  $OH^-$  and  $H_2O_2$  was not possible due to instrumental limitations.

Fully deuterated basic hydrogen peroxide (BHP) at  $-20^\circ\text{C}$  yielded a lifetime of  $20 \mu\text{s}$  for  $O_2(^1\Delta_g)$  by direct measurement, but a reliable direct measurement of  $\tau_{\Delta}$  under BHP conditions was not possible. However, using reasonable assumptions, we estimate  $\tau_{\Delta}$  in BHP at  $-20^\circ\text{C}$  to be  $1.8 \mu\text{s}$ .

The temperature dependencies of all the rate parameters have been measured, yielding activation energies close to  $6 \text{ kJ mol}^{-1}$ .

The current development of a novel singlet oxygen detector, which should have improved bandwidth, will be used to re-examine  $\tau_{\Delta}$  in BHP by direct measurement.



<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution/ _____	
<b>Availability Codes</b>	
Dist	Avail and/or Special
<b>A-1</b>	

## PREFACE

The Rotocoil singlet oxygen generator produces  $O_2(^1\Delta_g)$  chemically by reacting chlorine gas with basic hydrogen peroxide solution (BHP).

Although the lifetime of  $O_2(^1\Delta_g)$  has been measured in many solvent systems and has been found to be very solvent-dependent, it has never been determined under BHP conditions. In a solution of BHP such as that used in the Rotocoil generator, which is water with high concentrations of hydroxide ion ( $OH^-$ ) and hydrogen peroxide ( $H_2O_2$ ), the possibility of additional deactivation modes are introduced.

These deactivation processes under conditions of high concentrations of hydroxide and hydrogen peroxide have been investigated, as well as the effects of temperature on  $O_2(^1\Delta_g)$  deactivation.




## CONTENTS

Summary .....	iii
Preface .....	iv
1.0 Introduction .....	1
2.0 Experimental .....	5
2.1 Apparatus .....	
2.2 Sample Preparation .....	
3.0 Results .....	7
3.1. Direct Measurement of Singlet Oxygen Lifetime .....	
1 Perprotio-BHP .....	
2 Perdeutero-BHP .....	
3.2. Indirect Evaluation .....	
1 Quenching of $O_2(^1\Delta_g)$ by $D_2O_2$ .....	
2 Quenching of $O_2(^1\Delta_g)$ by $OD^-$ .....	
3 Quenching by $H_2O_2$ and $OH^-$ .....	
3.3 Temperature Dependence of $\tau_\Delta$ .....	
1 In $D_2O$ .....	
2 In 25wt% $D_2O_2$ in $D_2O$ .....	
3 In 3 M $OD^-$ in $D_2O$ .....	
4.0 Discussion .....	14
4.1 Comparison of Direct and Indirect Evaluations .....	
1 Perdeutero-BHP .....	
2 Estimation of Singlet Oxygen Lifetime in BHP .....	
5.0 Conclusions .....	16
References .....	17

## 1.0 INTRODUCTION

The lifetime of  $O_2(^1\Delta_g)$  in solution is solvent dependent (Refs. 1-4) and may vary by up to a factor of several hundred in common organic solvents. The lifetime ( $\tau_\Delta$ ) of the  $^1\Delta_g$  state also displays a solvent deuterium isotope effect (Ref. 1). In pure  $H_2O$ ,  $\tau_\Delta = 4 \mu s$ ; and in pure  $D_2O$ ,  $\tau_\Delta = 62 \mu s$ . The introduction of species such as  $OH^-$  and  $H_2O_2$  into the solvent system may introduce additional deactivation modes for the  $^1\Delta_g$  state (eqns. 6 and 7). Deactivation of the electronic excitation energy of  $O_2(^1\Delta_g)$  is thought to occur via energy transfer to solvent vibrational modes (Ref. 1).

The lifetimes of  $O_2(^1\Delta_g)$  in this report were determined by time-resolved studies of the singlet oxygen luminescence at 1270 nm ( $^1\Delta_g v=0$  to  $^3\Sigma_g v=0$  transition). Singlet oxygen was produced in solution by flash excitation of a sensitizing dye which undergoes energy transfer to oxygen from its triplet state. The overall kinetic scheme for production and decay is as follows:



The luminescence of singlet oxygen at 1270 nm was detected using a germanium-photodiode-based system. The decay of the  $O_2(^1\Delta_g)$  luminescence signal follows first-order kinetics. A representative decay trace is shown in Figure 1. A fast emission spike, thought to be due to sensitizer emission in the near infrared region, is seen at the beginning of the decay signal; however, after the decay of the spike, the kinetics are first-order. Semilogarithmic plots of the data are linear, Figure 2, confirming the first-order kinetics. The slope of the semilogarithmic plot yields the lifetime of singlet oxygen in that medium.






Figure 1. Luminescence decay of singlet oxygen at 1270 nm.




Figure 2. Semilogarithmic plot of figure 1.

However, the fast spike causes severe problems to measurements of systems having  $\tau_{\Delta}$  of a few microseconds. Luminescence yields from  $O_2(^1\Delta_g)$  are relatively weak compared to the intense fast emission. The latter has an exponential tail which, when  $\tau_{\Delta}$  are circa a few microseconds, merges with the singlet oxygen luminescence signal and becomes indistinguishable therefrom. For this reason we have been unable to obtain reliable direct data for the value of  $\tau_{\Delta}$  in BHP (protiated), see below.

## 2.0 EXPERIMENTAL

### 2.1 APPARATUS

The excitation source was a Quantel YG571C Q-switched Nd:YAG laser, producing circa 10 ns pulses with up to 1084 mJ per pulse at 1064 nm. The second (532 nm) and third (355 nm) harmonics were generated and separated from the fundamental by two successive dichroic mirrors and a heat-absorbing filter.


The luminescence from the irradiated solutions was detected using a germanium photodiode operating with a 6 volt reverse bias. The signal was fed into a Judson 100 preamplifier providing 55 dB of gain, followed by a differential amplifier operating at unit gain, to provide impedance matching with the input of the LeCroy 9400 digital oscilloscope. The LeCroy was on-line to an IBM PC/AT clone microcomputer. For all lifetime measurements, between 10 and 20 individual decay curves were taken and averaged in the computer. The entrance to the diode housing was covered with a 5-mm-thick piece of antireflection coated silicon which has a cut-on at 1050 nm. A 10x10 mm quartz fluorescence cell was placed in contact with the filter. The luminescence was monitored at 90° to the incident laser beam. The fluorescence cell was housed in an aluminium holder which could be heated or cooled from -30°C to +50°C by a circulating 1:1 water/ethylene glycol solution from a thermostatically controlled circulator. Temperature measurements were made using a Cole-Palmer 85671 bench indicator connected to a T-type thermocouple. This experimental setup is shown schematically in Figure 3.

### 2.2 SAMPLE PREPARATION

In aqueous solution the following sensitizers were used; rose bengal (RB), pyrene tetrasulfonic acid tetrasodium salt (PTSA) and meso-tetra (4-sulfonato-phenyl) porphine tetrasodium salt (TPPS). The optical densities at 532 nm and 355 nm were 0.6/cm. All solutions were oxygen saturated prior to irradiation. Saturation of the solutions with argon gas completely removed the  $O_2(^1\Delta_g)$  luminescence.

The solvents used were, D<sub>2</sub>O (99.8% atom), and deionized water purified through a nano-pure filter system, 30 wt% D<sub>2</sub>O<sub>2</sub> in D<sub>2</sub>O and 30 wt% H<sub>2</sub>O<sub>2</sub>.

Reagent grade potassium hydroxide and sodium deuterioxide (99.8% atom) were used as quenchers.



D - DICHOIC MIRROR  
 H - HEAT FILTER  
 S - SAMPLE CELL  
 F - SILICON FILTER  
 PA - PREAMPLIFIER  
 DA - DIFFERENTIAL AMPLIFIER  
 DO - DIGITAL OSCILLOSCOPE  
 PC - COMPUTER  
 OP - OUTPUT  
 PD - PHOTODIODE

Figure 3. Schematic of singlet oxygen luminescence detection system.

### 3.0 RESULTS

#### 3.1 DIRECT MEASUREMENTS OF THE SINGLET OXYGEN LIFETIME

##### 3.1.1 Perprotio BHP

For the reasons of luminescence impurity outlined above, a solution of BHP (25% H<sub>2</sub>O<sub>2</sub> and 25% KOH in H<sub>2</sub>O at -20°C) containing PTSA as sensitizer did not yield a reliable direct measurement of  $\tau_{\Delta}$ . It is possible to state that the value is not greater than 7  $\mu$ s. The indirect approach is detailed in section 3.2.

##### 3.1.2 Perdeutero BHP

Singlet oxygen deactivation by solvent shows a dramatic isotope effect. The enhancement achieved by fully replacing all H atoms by their deuterium counterpart are often in the range of 10-20 times the undeuteriated value. We prepared a sample of perdeutero BHP (BHP\*) containing 25% NaOD and 25% D<sub>2</sub>O<sub>2</sub> in D<sub>2</sub>O at -20°C with PTSA as sensitizer. Laser pulse excitation produced an infrared (IR) signal from singlet oxygen that was kinetically distinguishable from the decay of the rapid spike (Figure 4). From this we obtained  $\tau_{\Delta}(\text{BHP}^*) = 20 \mu\text{s}$ . This experiment could be performed only at sub-zero temperatures, as the BHP\* decomposed above -10°C.

To support this direct measurement and to provide data to allow an evaluation of  $\tau_{\Delta}(\text{BHP})$ , we carried out indirect measurements as detailed below.

#### 3.2 INDIRECT EVALUATION

The strategy of this approach is to evaluate bimolecular rate constants (where possible) for the deactivation processes:



and the temperature dependences of these processes. In this way we can expect to evaluate  $\tau_{\Delta}$  from the relationship:

$$\tau_{\Delta}(\text{BHP})^{-1} = k_d(\text{BHP}) = k_{\text{O}} + k_{\text{OH}^-}[\text{OH}^-] + k_{\text{H}_2\text{O}_2}[\text{H}_2\text{O}_2] \quad (12)$$




Figure 4. Luminescence decay of singlet oxygen in deuterated BHP.

where  $k_O = (\tau_\Delta)^{-1}$  is the deactivation rate constant in neat water. These  $k_O$  values for both  $H_2O$  and  $D_2O$  are known from the literature (Ref. 4).

### 3.2.1 Quenching of $O_2(^1\Delta_g)$ by $D_2O_2$

$D_2O_2$  was shown to quench  $O_2(^1\Delta_g)$  only weakly. Figure 5 depicts a plot of the decay rate ( $s^{-1}$ ) of the IR luminescence as a function of  $D_2O_2$  concentration in  $D_2O$  over the range 0-10 M at 20°C. The plot is linear as expected for a first-order dependence on concentration. The bimolecular rate constant ( $k_{D_2O_2}$ ) evaluated from the slope of Figure 5 is  $8.8 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$ .

### 3.2.2 Quenching of $O_2(^1\Delta_g)$ by $OD^-$

An analogous series of experiments to those described in 3.2.1 were carried out at 20°C but replacing  $D_2O_2$  by  $NaOD$ . The decay of  $O_2(^1\Delta_g)$  luminescence was exponential and Figure 6 shows a linear relationship between the measured decay constant and  $OD^-$  concentration. From this  $k_{OD^-} = 5.5 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ .

### 3.2.3 Quenching by $H_2O_2$ and $OH^-$

The lifetime of singlet oxygen in  $H_2O$  is 4  $\mu s$ , which is at the low end of reliable measurements because of sensitizer-derived emission spilling over in the IR measuring region. Thus we were unable to measure any decreases in this value caused by quenching additives such as  $H_2O_2$  and  $OH^-$ .

## 3.3 TEMPERATURE DEPENDENCE OF $\tau_\Delta$

### 3.3.1 IN $D_2O$

Over the temperature range 278-323 K, the value of  $\tau_\Delta(D_2O)$  varied from 71  $\mu s$  to 48  $\mu s$  with a value of 61  $\mu s$  at room temperature (20°C). An Arrhenius plot of the data using a variety of sensitizers (Figure 7) showed linear behavior from which the relationship (equation 13) is obtained.

$$k_O(T) = 1.88 \times 10^5 \exp(-714/T) \text{ s}^{-1} \quad (13)$$

### 3.3.2 In 25% $D_2O_2$ in $D_2O$

For a 25%  $D_2O_2$  solution in  $D_2O$  measured over a range of temperature from 258-283 K, an Arrhenius plot of the first-order deactivation rate parameter (in  $s^{-1}$ ) yielded

$$k_d(D_2O_2/D_2O) = 3.57 \times 10^6 \exp(-1408/T) \text{ s}^{-1} \quad (14)$$






Figure 5.  $k_d$  vs concentration of deuterium peroxide.




Figure 6.  $k_d$  vs concentration NaOD.




Figure 7. Arrhenius plot of temperature dependence of singlet oxygen lifetime.

### 3.3.3 3 Molar OD<sup>-</sup> in D<sub>2</sub>O

An Arrhenius plot of the first-order decay of O<sub>2</sub>(<sup>1</sup>Δ<sub>g</sub>) in an alkaline (OD<sup>-</sup>) solution in D<sub>2</sub>O yielded

$$k_d(\text{OD}^-/\text{D}_2\text{O}) = 3.8 \times 10^5 \exp(-660/T) \text{ s}^{-1} \quad (15)$$

## 4.0 DISCUSSION

### 4.1 COMPARISON OF DIRECT AND INDIRECT EVALUATIONS

#### 4.1.1 Perdeutero-BHP

As indicated before, the deactivation rate constant of a mixture of BHP components in the general case is given by:

$$k_T = k_O + k_p[\text{peroxide}] + k_g[\text{base}] \quad (16)$$

where  $k_O$  is the first-order constant for the deactivation in the neat solvent, and  $k_p$  and  $k_g$  are bimolecular rate constants for deactivation by peroxide and base species respectively. All rate parameters in equation (16) are temperature dependent, but only very weakly so ( $E_a$  values are near 6 kJ mol<sup>-1</sup>). For deuterated materials, we have determined values of  $k_O$ ,  $k_p$  and  $k_g$  at 20°C. These are 1.7x10<sup>4</sup> s<sup>-1</sup>, 8.8x10<sup>2</sup> M<sup>-1</sup>s<sup>-1</sup>, and 5.5x10<sup>3</sup> M<sup>-1</sup>s<sup>-1</sup>, respectively.

The BHP and BHP\* systems we made contained 25% by weight of peroxide and base components. This yields [OD<sup>-</sup>] = 4.8 M and [D<sub>2</sub>O<sub>2</sub>] = 6.6 M. From these rate constants and concentration values we can evaluate  $k_T$  (BHP\*) from

$$k_T = 1.7 \times 10^4 + (8.8 \times 10^2 \times 6.6) + (5.5 \times 10^3 \times 4.8) = 4.9 \times 10^4 \text{ s}^{-1} \quad (17)$$

and, since  $\tau_\Delta(\text{BHP}) = k_T(\text{BHP}^*)^{-1}$ ,  $\tau_\Delta(\text{BHP}^*) = 20.4 \text{ } \mu\text{s}$  at 20°C by indirect evaluation, which compares to 20  $\mu\text{s}$  by direct measurement at -20°C. Thus, from  $k_O = 1.88 \times 10^5 \exp(-714/T) \text{ s}^{-1}$ , we find  $k_O(-20^\circ\text{C}) = 1.13 \times 10^4 \text{ s}^{-1}$ .

From  $k_d(T) = 3.8 \times 10^5 \exp(-880/T) \text{ s}^{-1}$  for 3 M OD<sup>-</sup>/D<sub>2</sub>O, we find  $k_g(\text{OD}^-)(-20^\circ\text{C}) = 2.7 \times 10^4 \text{ s}^{-1}$  at 4.8 M OD<sup>-</sup>.

From  $k_d(T) = 3.57 \times 10^6 \exp(-1408/T) \text{ s}^{-1}$  for 6.6 M D<sub>2</sub>O<sub>2</sub>/D<sub>2</sub>O, we find  $k_p[\text{D}_2\text{O}_2] = 3.0 \times 10^3 \text{ s}^{-1}$  at 6.6 M D<sub>2</sub>O<sub>2</sub>.

Therefore  $\Sigma k = 1.13 \times 10^4 + 2.7 \times 10^4 + 3.0 \times 10^3 = 4.1 \times 10^4 \text{ s}^{-1}$  and  $\tau_\Delta = \Sigma k^{-1} = 24.3 \text{ } \mu\text{s}$ .

This value is close enough to the direct measurement of 20  $\mu\text{s}$  at -20°C for  $\tau_\Delta$  in BHP\* to give us confidence in the procedure. One useful fact to

note is that the contribution of  $D_2O_2$  to the total decay is only circa 7% and is therefore relatively insignificant.

#### 4.1.2 Estimation of Singlet Oxygen Lifetime in BHP

Experimental problems have prevented the direct measurement of  $\tau_\Delta$  in BHP and the values of  $k_{OH^\cdot}$  and  $k_{H_2O_2}$  in order to make the indirect evaluation. However, an estimate can be made as follows.

In the equation for overall decay:

$$k_T = k_O + k_B[\text{Base}] + k_P[\text{Peroxide}]$$

We have shown in the previous section that the  $k_P[\text{Peroxide}]$  factor is small enough to be ignored. Then

$$k_T = k_O + k_B[\text{Base}] \quad (18)$$

We know  $k_O$  for  $H_2O$  at  $20^\circ C$  from the literature, and by assuming the same exponential temperature parameter (714 K) as for  $D_2O$ , we can equate the ratios of the pre-exponential factors (A values) to that of the rate constant at  $20^\circ C$ .

$$\text{Thus:} \quad A_O(H) = A_O(D) \cdot k_O(H)/k_O(D) \quad (19)$$

$$= \frac{1.88 \times 10^5 \times 2.5 \times 10^5}{1.7 \times 10^4} = 2.8 \times 10^6 \text{ s}^{-1}$$

From this we write down the Arrhenius expression for  $k_O(H_2O)$  as

$$k_O(H_2O) = 2.8 \times 10^6 \exp(-714/T) \text{ s}^{-1} \quad (20)$$

and, at  $-20^\circ C$  (253 K), this will contribute  $1.7 \times 10^5 \text{ s}^{-1}$  to the total decay ( $k_T$ ).

We cannot use the same expression to estimate  $k(OH^\cdot)$  since we do not have data for  $OH^\cdot/H_2O$  solution at  $20^\circ C$ . However, we can make an argument based upon literature values for relative contributions ( $\Delta_{ox}$ ) to the singlet oxygen deactivation by OH and OD oscillators in alkanols (Ref. 5). Thus,  $\Delta_{OH} = 2290$  and  $\Delta_{OD} = 165$ ; and if we assume the ratio (H/D = 14) (Ref. 5) applies equally well to  $OH^\cdot$  and  $OD^\cdot$  species, then we can convert  $k_{OD^\cdot}[OD^\cdot] = 2.7 \times 10^4 \text{ s}^{-1}$  at  $-20^\circ C$  into  $k_{OH^\cdot}[OH^\cdot] = 3.8 \times 10^5 \text{ s}^{-1}$ .

In equation (18), therefore,

$$k_T = 1.7 \times 10^5 + 3.8 \times 10^5 = 5.5 \times 10^5 \text{ s}^{-1} \text{ at } -20^\circ C$$

or  $\tau_\Delta(\text{BHP}) = 1.8 \text{ } \mu\text{s}$  in BHP at  $-20^\circ C$ .

## 5.0 CONCLUSIONS

These experiments have led to the following conclusions:

- (1) Fully deuterated BHP at  $-20^{\circ}\text{C}$  supports a lifetime of  $20\ \mu\text{s}$  for  $\text{O}_2(^1\Delta_g)$  by direct measurement. This compares to  $60\ \mu\text{s}$  at room temperature for  $\text{D}_2\text{O}$  alone.
- (2) The contributing bimolecular rate constants  $k_{\text{OD}^{\cdot}}$  and  $k_{\text{D}_2\text{O}_2}$  have been determined and their temperature dependence measured. These data have allowed an indirect estimation of  $\tau_{\Delta}$  for deuterio-BHP of  $24\ \mu\text{s}$ , very close to the directly measured value.
- (3) Instrumental limitation prevented direct measurement of  $\tau_{\Delta}$  in BHP at  $-20^{\circ}\text{C}$  and of  $k_{\text{OH}^{\cdot}}$  and  $k_{\text{H}_2\text{O}_2}$ . However, using reasonable assumptions, we estimate  $\tau_{\Delta}$  in BHP at  $-20^{\circ}\text{C}$  to be  $1.8\ \mu\text{s}$ .
- (4) After completion of this program, a novel singlet oxygen detector has been developed which should have improved bandwidth. If so, it will be used to re-examine BHP (direct).

## REFERENCES

1. P. R. Ogilby and C. S. Foote; " Effect of Solvent, Solvent Isotopic Substitution, and Temperature on the Lifetime of Singlet Molecular Oxygen ( $^1\Delta_g$ ). " J. Amer. Chem. Soc., 105, 3423, 1983.
2. M. A. J. Rodgers ; " Time Resolved Studies of 1.27  $\mu\text{m}$  Luminescence from Singlet Oxygen Generated in Heterogenous and Microheterogenous Fluids.", Photochem. Photobiol., 37, 99, 1983.
3. C. A. Long and D. R. Kearns; "Radiationless Decay of Singlet Molecular Oxygen in Solution. 11. Temperature Dependence and Solvent Effects ." J. Amer. Chem. Soc., 97, 2018, 1974.
4. M. A. J. Rodgers and P. T. Snowden, " Lifetime of  $\text{O}_2(^1\Delta_g)$  in Liquid Water as Determined by Time-Resolved Infrared luminescence Measurements.", J. Amer. Chem. Soc., 104, 5541, 1982.
5. M. A. J. Rodgers, "Solvent-Induced Deactivation of Singlet Oxygen : Additivity Relationships in Nonaromatic Solvents.", J. Amer. Chem. Soc. , 105, 6201, 1983.

## GENERAL REFERENCES

1. W. C. Schumb, C. N. Satterfield and R. L. Wentworth, " Hydrogen Peroxide ", Reinhold, New York, 1955.
2. " Singlet  $\text{O}_2$  ", Ed., A. A. Frimer, CRC Press, Florida, 1985.